

IMPACT OF OPEN SUB SURFACE DRAINAGE ON RECLAMATION OF WATERLOGGED AND SALT AFFECTED SOILS

G. KISHORE BABU¹, S.RAMESH CHANDRA², K.YELLA REDDY³, G.SUBBA RAO⁴
AND Y.RADHA⁵, A.P. WATER MANAGEMENT PROJECT, BAPATLA-522101

ABSTRACT

Soil, land and water are essential resources for the sustained quality of human life and foundation of agricultural development. Efficient management of land and water resources is a major challenge for the scientist, planners, administrators and farmers to ensure food, water and environmental security for the present and future generations (Kanwar, 2000, Das, 2009). Irrigation development is largely responsible for making India self sufficient in food grains production but the negative aspect is drainage has not been given importance as much as irrigation. Productive agricultural land is going out of cultivation because water logging and salinity caused by the raise of water table due lack of proper drainage are the major problem in the canal command areas. Adoption of Sub surface drainage technology is probably one of the best ways to increase resource use efficiency in order to increase crop production and sustain natural resources like soil and water in severely water logged saline soils. So there is a great demand for the research and developmental efforts to reclaim all the salt affected and water logged soils by providing drainage and bring them back to non saline productive soils.

The Andhra Pradesh Water Management Project has conducted detailed study in the farmers fields of drainage pilot area located in Pathareddy palem village under Mutluru channel command of Krishna Western Delta, Guntur District to evaluate the performance of Open Subsurface Drainage system (OSSD) for improving the grain yield of Rice under saline and water logged conditions for three consecutive seasons starting from Kharif, 2005. The grain yield was increased from 2.2 t/ha to 4.9t/ha over a period of four years indicating an over all increase of 123% due to installation of OSSD. The cropping intensity and non saline area were also increased to 100-158% and 6-40% respectively.

The yields of major food grains have stagnated or declined since 2000. One of the ways to improve the food grain production is, restoring degraded or problematic soils and improving quality of surface and ground water (Lal, 2008). The total salt-affected area in the state of Andhra Pradesh is estimated as 8.18 lakh ha. The irrigation induced water logging and salinity were estimated to be 2.72 and 1.15 lakh ha respectively and these are the two major causes which lead to reduction in crop yields in most of the irrigated commands (Anonymous, 2006). It is also observed that continuous irrigation over the years without proper drainage also results in raising the ground water table to the root zone, thus leading to reduction in crop yields. From the results of the Reconnaissance Survey, an area of 35000 ha was identified as salt affected in the Krishna Western Delta (KWD) irrigated commands where the average grain yield of rice was as low as 1.5-2.5t/ha.

The problem of increasing salinity caused by the rise of water table due to lack of proper drainage is considered as a major environmental problem that threatens the capital investment in irrigated agriculture and its sustainability. Drainage has not been given importance as much as irrigation by the farmers as well as the Government Agencies. So there is a great demand for the concerned research and development efforts to reclaim all the salt affected and water logged soils and bring them back to profitable farming with increased agricultural production as well as cropping intensity. The only means to overcome the salinity and water logging permanently is selection and adoption of suitable Sub Surface Drainage systems. To test and demonstrate the drainage need for control of soil salinity and water logging, a collaborative project of Acharya N.G.Agricultural University with Alterra, ILRI, The Netherlands, was approved by the Government of Andhra Pradesh Bapatla as a main center. The Bapatla center is entrusted with the responsibility of solving the water logging and salinity problems by installation of open subsurface drainage (OSSD) systems in the canal commands of Andhra Pradesh.

¹Senior Scientist (Soil Science), ²Scientist (SWE), ³Principal Scientist & Project Manager, ⁴Senior Scientist (Agronomy) and ⁵Senior Scientist (Agril. Economics), A.P Water Management Project, Bapatla 522 101

MATERIALS AND METHODS

A pilot area of 22.4 ha was selected at Pathareddy palem village under Mutluru channel command of KWD in Guntur district to study the performance of open sub surface drainage system to overcome the severe problem of salinity and water logging.

. The soil samples collected from 62 grid points of the pilot area (Fig1) fixed at spacing of 60mts apart were analyzed for pH, ECe, water-soluble cations and anions, ESP, and available macro and micronutrients following the standard procedures described by Black (1965) and Richards (1954). Crop cutting experiments have also been conducted at each of these grid points to estimate the crop yields during 2005-08.




Fig. 1 Map showing the grid soil sample locations in Open sub surface drainage system at Pathareddy Palem drainage pilot area

The soils of the pilot area are deep and clayey in texture, with ECe ranging from 1.6 to 61.3 dS/m and pH from 6.7 to 8.4. Sodium is the dominant water soluble cation followed by Mg and Ca, while chlorides are the dominant water soluble anions followed by sulphates and bicarbonates. The ESP varies from 10.12 to 14.10 indicating that the soils are saline and not sodic in nature. The available nitrogen is low to medium, while the phosphorous and potassium are medium to high. The average values of ECe of ground water are highly saline and not suitable for agriculture. The soils are highly affected by water logging and salinity due to insufficient capacity of main drain to evacuate excess rainfall. The principal crop grown in the pilot area is rice during *Kharif* season.

The open sub surface drainage system (Fig 2) consists of 10 laterals covering a length of 1250 m and bottom width of 0.75 m with slope of 0.01% gradient, being properly dugged out with hydraulic excavator. The laterals were first connected to a collector drain. The

collector drain was connected to a secondary drain and then to the Mutluru side drain to convey drain water offsite. The drainage coefficient was 1mm/day. Eight observation wells were installed up to 3.4 m depth in Mutluru channel command and four out of eight falls in the experimental site to monitor water table fluctuations and for sampling ground water for water quality determinations.




Fig. 2 Map showing Open sub surface drainage system at Pathereddy Palem drainage Pilot area under Mutluru Channel Command of Krishna Western Delta

RESULTS AND DISCUSSION

Monitoring of the open subsurface drainage system during *Kharif* 2005-2008 clearly indicated that the system effectively controlled the ground water table in the root zone at a relatively shallow depth of 0.5-1.5mts. The shallow water table also avoids excessive drainage while at the same time harmful salts that are brought in by the irrigation water were also reduced to a level ranging from 1.2 to 35.4 dS/m. Further the data on soil salinity at 62 grid locations showed a relative decrease in soil salinity during post drainage conditions with an average mean value of 11.6, 14.2, 12.9 and 11.8 dS/m (Table 1) during first, second, third and fourth year after installation respectively, which helped in increasing the grain yield. Prasad *et al.*, (2007) also reported positive results with open sub surface drainage system in reducing the salinity of problematic soils.

The grain yield (Table 1, Fig. 4) data collected from 62 grid points indicated a steady increase from 2005 to 2008. The average grain yield obtained during *Kharif* 2005, 2006, 2007 and 2008 were 2.9 t/ha, 3.8 t/ha, 4.5 t/ha and 4.9 t/ha respectively, as against 2.2 t/ha (2004) before installation of the system. Further, an overall increase of 123 per cent in grain yield was observed by the end of four years after installation of the drainage system. Similar

positive results were reported by Konanki and Uppugundur drainage pilot areas of Indo-Dutch net work project. (IDNP Comprehensive Report, 1995)

The cropping intensity was 100% with *Kharif* rice only without any second crop before installation of the system where as the cropping intensity was also increased to 158 % by the end of four year after installation of the system allowing the farmers to raise Fodder Jowar, sunhemp and black gram as second crop during *rabi* season in 58% of the drainage pilot area. The significant increase in crop yield and cropping intensity can be attributed to the direct effects of the introduction of open subsurface drainage system which in turn lowered the water table and decreased the soil salinity by leaching out the soluble salts from the root zone, there by not only creating the favorable conditions in root zone but also making the nutrients available to the plants resulting in optimum plant growth and yield.

Further, the shallow water table helped to overcome the excessive drainage and also remove the harmful salts that are brought in by the irrigation water there by creating favorable conditions in the soil root zone to establish fodder crops successfully in the second season after harvest of *Kharif* rice.

The operational research conducted in the farmer's fields of drainage pilot area at Pathareddy palem indicated that, under the prevailing soil and agro climatic conditions Open subsurface drainage system is technically feasible to reclaim water logged and saline soils and to sustain agricultural production in irrigated commands. Further, the Open Sub-surface Drainage System needs frequent de-silting of drains during summer months for effective functioning of the system though it requires a huge amount of labor work. Thus, Open Subsurface Drainage system is one of the best tools for permanent reclamation of water logged and saline soils in Krishna Western Delta Commands




Fig.3 Impact of Open sub-surface drainage system on summer soil salinity of Pathareddy Palem pilot area

Table1. Effect of Open Sub-Surface Drainage System on soil salinity, grain yield and cropping intensity during 2005-2008

Year	ECe dSm ⁻¹	Grain Yield (t ha ⁻¹)	Cropping intensity (%)
2004 (before installation)	21.7	2.2	100
2005	11.6	2.9	125
2006	14.2	3.8	146
2007	12.9	4.5	153
2008	11.8	4.9	158




Fig. 4 Relationship between ECe and Grain yield during 2005-2008

LITERATURE CITED

Annual reports 2002-2006 All India Co-Ordinated Project on management of salt affected soils and use of saline water in Agriculture.

Black C. A. 1965 Methods of Soil Analysis, Part-1 & II American Society of Agronomy Madison, Wisconsin, USA.

Comprehensive Report on Research Achievements of Indo-Dutch Net Work Project, Bapatla (1-4-96 to 31-3-2002) on “Drainage and Water Management for Salinity Control in Canal Commands”.

Das D.K, Bandyopadhyay S, Chakrabarty D and Rajeev Srivatava 2009 Application of modern techniques in characterization and management of soil and water resources. Journal of Indian Society of Soil Science vol. 57 (4) pp 445-460.

Kanwar J S 2000 Soil and water resources management for sustainable agriculture-imperatives for India. In. Development and conservation, Int. Conf. on managing natural resources for sustainable agricultural production in 21st century. Feb 14-18, New Delhi pp 17-37

Lal R. 2008 Soils and India's food security. Journal of Indian Society of Soil Science, 56(2):129-138

Prasad P. R. K., Srinivas D, T. V. Satyanarayana, Ramesh Chandra S., Subba Rao G., Mukunda Rao B. and Srinivasulu A. 2007. Reclamation saline and waterlogged soils in Mutluru channel command of Krishna Western Delta, Andhra Pradesh state, India Paper presented during 58th ICID meeting and The USCID 4th International conference on irrigation and drainage organized from 30-09-2007 to 06-10-2007 at Sacramento Convention Center, California-USA.

Richards L. A. 1954. Diagnosis and Improvement of Saline and Alkali Soils. USDA Handbook 60, US Department of Agriculture and Oxford & IBH Pub. Co. Pvt. Ltd., New Delhi.